

LA-UR-14-23050

Approved for public release; distribution is unlimited.

Title: Ultracold Neutrons at LANSCE

Author(s): Broussard, Leah J.

Intended for: PILLAR Institute for Lifelong Learning Tour Group

Issued: 2014-04-30



Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.


Ultracold Neutrons at LANSCE

Leah Broussard

Los Alamos National Laboratory

February 29, 2014

The Standard Model (and Beyond)




Some Curiosities

- Lots of “Why?’s”
 - Why 3 generations?
 - Why so many parameters?
 - Why these masses?
 - Why left-handed weak interaction?
- What is Dark Matter?
- Why so much matter?
- Where is gravity?
- And more. . .

Finding the missing pieces

- High Energy frontier (LHC) vs. Precision frontier (beta decay)
- Complementary approaches
 - High energy: Direct search for heavy particles
 - Precision: Measure deviations from SM expectation

Neutron Beta Decay



- Semileptonic charged weak interaction
- Standard Model: **V** - **A** (left-handed)
- Lifetime ~ 15 minutes

What we can measure:

- Total decay rate (lifetime):

$$\frac{1}{\tau_n} = W = K (G_F \mathbf{V}_{ud})^2 \left(1 + 3 \left(\frac{G_A}{G_V} \right)^2 \right) (1 + \Delta_R) f_n p_e E_e (E_0 - E_e)^2 \left[1 + m_e \mathbf{b} \frac{f_b}{f_n} \right]$$

- Angular correlations:

$$\frac{dW}{dE_e d\Omega_e d\Omega_\nu} \propto p_e E_e (E_0 - E_e)^2 \left[1 + \mathbf{a} \frac{\vec{p}_e \cdot \vec{p}_\nu}{E_e E_\nu} + \mathbf{b} \frac{m_e}{E_e} + \langle \vec{\sigma}_n \rangle \cdot \left(\mathbf{A} \frac{\vec{p}_e}{E_e} + \mathbf{B} \frac{\vec{p}_\nu}{E_\nu} + \mathbf{D} \frac{\vec{p}_e \times \vec{p}_\nu}{E_e E_\nu} \right) \right]$$

Testing the Standard Model


- **A**, **a** + $\tau \rightarrow$ **V**, **A** interactions (\mathbf{V}_{ud} , RL symmetry, ...)
- **B**, **b** \rightarrow **S**, **T** interactions (BSM interactions, MSSM, ...)

Ultracold Neutrons

Class	Energy	Source
Fast	$> 1 \text{ MeV}$	Fission reactions / Spallation
Slow	$\text{eV} - \text{keV}$	Moderation
Thermal	0.025 eV	Thermal equilibrium
Cold	$\mu\text{eV} - \text{meV}$	Cold moderation
Ultracold	$\leq 300 \text{ neV}$	Downscattering

How cold is Ultracold?

- Temperature $< 4 \text{ mK}$
- Velocity $< 8 \text{ m/s}$
- Usain Bolt $\sim 12 \text{ m/s}$



UCN can be bottled

- Gravitational ($V = mgh$): $100 \text{ neV} / \text{meter}$
- Magnetic ($V = -\vec{\mu} \cdot \vec{B}$): $60 \text{ neV} / \text{Tesla}$
- Material $\left(V = \frac{2\pi\hbar^2 Nb}{m} \right) \left\{ \begin{array}{ll} {}^{58}\text{Ni} : & 335 \text{ neV} \\ \text{DLC} : & 250 \text{ neV} \\ \text{BeO} : & 250 \text{ neV} \\ \text{Cu} : & 170 \text{ neV} \end{array} \right.$

Manipulating UCN


Material bottles: UCN guides

- Features of good UCN guides:
 - Low “loss per bounce” ($< 10^{-5}$)
 - High Material Potential (> 200 neV)
 - Low depolarization ($\sim 10^{-6}$)




Magnetic selection: UCN polarizers

- Neutron magnetic moment μ due to spin
- 100% polarization possible





(note: neutron magnetic moment is negative)



Ultracold Neutron Facility at LANSCE





- 800 MeV proton beam + tungsten target → spallation neutrons
- Single scatter in solid deuterium: $CN \rightarrow UCN + \text{phonon}$
- Remove phonons: SD_2 cooled to 4K
- “Flapper” shields UCN from SD_2
- 50 UCN/cc at shield wall
- Pulsed beam: Low background



Experimental Programs at the UCN Facility: UCN^A




Experimental Programs at the UCN Facility: UCN**A**





Corr. +/- Uncertainty (%)	Mendenhall (2013)	In analysis (TBS 8/14)	Next Step	Source of improvement
Statistics	+/- 0.46	+/- 0.40	+/- 0.28	Decay rate!
Depolarization	+0.67 +/- 0.56	+0.67 +/- 0.1	+0.6 +/- 0.1	Shutter+ ex situ
Backscatter	+1.36 +/- 0.34	+0.5 +/- 0.15	+0.5 +/- 0.15	Thin windows
Angle effect	-1.21 +/- 0.30	-0.8 +/- 0.2	-0.8 +/- 0.1	Windows+ APD
Energy Reconstruction	+/- 0.31	+/- 0.08	+/- 0.08	Xenon + LED
Total Sys.	+/- 0.82	+/- 0.28	+/- 0.22	
Total	+/- 0.94	+/- 0.5	+/- 0.35	
	Dominated by systematics	Dominated by statistics	Intended to be balanced	


Experimental Programs at the UCN Facility: UCN**B**



- 3-body decay: ν -asymmetry from proton/electron directions
- **B** is sensitive to **b**, $\mathbf{b}_\nu \rightarrow$ non-Standard Model Scalar and Tensor interactions



Experimental Programs at the UCN Facility: UCNB



Experimental Programs at the UCN Facility: UCNB



Novel thick, large area, highly segmented silicon detectors



Biased to -30 kV to accelerate protons



Custom preamplifiers: 24 ch instrumented




Experimental Programs at the UCN Facility: UCN τ

Neutron lifetime

- Precise tests of Standard Model
- Big bang nucleosynthesis
- Reactor and solar neutrino flux predictions

Beam vs. Bottle




- Beam experiments: neutron flux?
- UCN bottle experiments: loss in the walls?



Experimental Programs at the UCN Facility: UCN τ


Magneto-gravitational Trap

- World's largest permanent magnet array
- Neutrons bounded by magnetic field on bottom, gravity on top
- No wall losses!



Experimental Programs at the UCN Facility: UCN τ

First storage time measurement (Feb 2013)



Current status

- Precision studies of systematics
- In-situ UCN counting
- Quasibound UCN cleaning?
- Monte Carlo UCN simulations
- Goal for prototype: 1 s precision measurement
- Ultimate experimental goal: 0.1 s precision